

HER
1237**Identification**

<i>Genus</i> Halobacterium	<i>Species</i> salinarum	<i>Subspecies</i>	<i>Strain</i> NRC 34001
-------------------------------	-----------------------------	-------------------	----------------------------

*Aliases***Growth conditions**

<i>Temp.</i> 37	<i>Medium</i> NaCl?	<i>Aerobic/Anaerobic</i>	<i>Agitation</i>	<i>Other</i>
--------------------	------------------------	--------------------------	------------------	--------------

Lysogeny**Phage sensitivity**

HER 237 (S45)

HER 343 (Ja.1)

Reference

Lockhead, A.G. 1934. Can. J. Res. 10: 275-286.

Remark

Also held in the NRC permanent collection (no. 34001) and in the ATCC collection (no. 33170)
For reproducible results, it is best to maintain the organism in log phase in broth, checking frequently that the cells demonstrate regular morphology and appear as slender rods.
Log phase cultures can be refrigerated for 2-3 weeks.

History

History

Received from

Dr Allen C. Wais, Department of Biochemistry, Faculty of
Medecine, U.W.I. Mona, Kingston 7, Jamaica, West Indies

Date

3-22-1985

Received from

R.E. MacDonald, Cornell University, Ithaca, NY, USA

Date

02-1973

Received from

D.J. Kushner, Dept. Of Biology, University of Ottawa

Date

Last update

2012-02-24