

*HER*  
1307**Identification**

<i>Genus</i> Vibrio	<i>Species</i> parahaemolyticus	<i>Subspecies</i>	<i>Strain</i> 16
------------------------	------------------------------------	-------------------	---------------------

*Aliases***Growth conditions**

<i>Temp.</i> 25	<i>Medium</i> NaCl	<i>Aerobic/Anaerobic</i>	<i>Agitation</i>	<i>Other</i>
--------------------	-----------------------	--------------------------	------------------	--------------

**Lysogeny****Phage sensitivity**

HER 307 (?16)	HER 308 (?HAWI-5)	HER 309 (?PEL8C-1)
---------------	-------------------	--------------------

*Reference*

Kellogg, C.A., J.B. Rose, S.C. Jiang, J.M. Thurmond, and J.H. Paul. 1995. Genetic diversity of related vibriophages isolated from marine environments around Florida and Hawaii, USA. Marine Ecol. Progr. Ser. 120:89-98.

*Remark*

Some natural resistance to nalidixic acid.  
Has photoreactivation and nucleotide excision repair of UV damage.  
Used to isolate over 200 phages from marine environments around Florida, the Gulf of Mexico, Hawaii.

**History***History***Received from**

Dr Christina A. Kellogg, University of South Florida, Marine Science Department, 140 7th Avenue S., St. Petersburg, FL33701, USA

**Date**

09-22-1998

*Last update*

2012-02-29