


<i>HER number</i> 135			
Identification			
<i>Name</i> 135	<i>Morphotype</i> C1 (Podophage)	<i>Other designations</i>	
Taxonomy			
<i>Realm</i> <i>Duplodnaviria</i>	<i>Kingdom</i> <i>Heunggongvirae</i>	<i>Phylum</i> <u><i>Uroviricota</i></u>	<i>Class</i> <i>Caudoviricetes</i>
<i>Order</i>	<i>Family</i>	<i>Genus</i> <i>Perisivirus</i>	<i>Species</i> <i>Perisivirus Pr</i>
Images			
<i>Electron Micrograph</i> <i>Image</i> 	<i>Image description</i> Magnification: 297,000X Bar: 50 nm Staining: PTB		

<p><i>Characteristics</i></p> <p>Plaques: 0.2- 0.5mm, clear, sloping borders. Host is class 3 pathogen. Lytic also for <i>*B. ovis*</i> and rough strains of <i>*B. abortus*</i>. Grows well on <i>*B. canis*</i> RM6/66.</p>	<p><i>Genomic sequence</i> Activated</p>
<p>Propagation conditions</p>	
<p><i>Bacterial hosts</i> 1135</p>	
<p><i>Reference</i> Ackermann, H.-W., F. Simon, and J.-M. Verger. A survey of <i>*Brucella*</i> phage and morphology of new isolates. <i>Intervirology</i>. 1981. 16:1-7.</p>	
<p><i>Remarks</i></p>	
<p>History</p>	
<p><i>History</i></p> <p>Received from M. J. Corbel, FAO/WHO Collaborating Centre for Reference and Research on Brucellosis, Ministry of Agriculture, Fisheries and Food, Central Veterinary Laboratory, New Haw, Weybridge, Surrey, England, KT15 3NB</p>	<p>Date 09-28-1982</p>
<p><i>Source</i> Derived from R/O phage, England.</p>	
<p><i>Updated at</i> 2024-01-16</p>	