


<i>HER number</i> 152			
Identification			
<i>Name</i> 152	<i>Morphotype</i> B2 (Siphophage)	<i>Other designations</i>	
Taxonomy			
<i>Realm</i> <i>Duplodnaviria</i>	<i>Kingdom</i> <i>Heunggongvirae</i>	<i>Phylum</i> <u><i>Uroviricota</i></u>	<i>Class</i> <i>Caudoviricetes</i>
<i>Order</i>	<i>Family</i>	<i>Genus</i>	<i>Species</i>
Images			
<i>Electron Micrograph</i> <i>Image</i>  H.-W. ACKERMANN	<i>Image description</i> Magnification: 297,000X Bar: 50 nm Staining: UA		

<p><i>Characteristics</i></p> <p>Plaques: 0.2mm, clear. Sensitive to lyophilisation. Read plates after 48h - 72 hrs on agar.</p>	<p><i>Genomic sequence</i> Deactivated</p>
<h2>Propagation conditions</h2>	
<p><i>Bacterial hosts</i> 1152</p>	
<p><i>Reference</i> Patel, J. J. 1976. Morphology and host range of virulent phages of lotus rhizobia. Can. J. Microbiol. 22: 204-212.</p>	
<p><i>Remarks</i></p>	
<h2>History</h2>	
<p><i>History</i></p> <p>Received from Dr J. J. Patel, Department of Scientific and Industrial Research, Applied Biochemistry Division, Private Bag, Palmerston North, New Zealand.</p>	<p>Date 11-01-1982</p>
<p>Isolated by J. J. Patel From:</p>	<p>Date 1975 ?</p>
<p><i>Source</i> Soil, Palmerston North, New Zealand</p>	
<p><i>Updated at</i> 2024-01-16</p>	