

<i>HER number</i> 167			
Identification			
<i>Name</i> 167	<i>Morphotype</i> B2 (Siphophage)	<i>Other designations</i>	
Taxonomy			
<i>Realm</i> <i>Duplodnaviria</i>	<i>Kingdom</i> <i>Heunggongvirae</i>	<i>Phylum</i> <u><i>Uroviricota</i></u>	<i>Class</i> <i>Caudoviricetes</i>
<i>Order</i>	<i>Family</i>	<i>Genus</i>	<i>Species</i>
Images			
<i>Electron Micrograph</i> <i>Image</i>  FHRCBV H.-W. ACKERMANN	<i>Image description</i> Magnification: 297,000X Bar: 50 nm Staining: UAB		
<i>Characteristics</i> Plaques of 1-1.5 mm, target-type. Elongated head.	<i>Genomic sequence</i> Deactivated		

Propagation conditions

Bacterial hosts

1165

Reference

Koga, T., S. Toyoshima, and T. Kawata. 1982. Morphological varieties and host ranges of **Vibrio parahaemolyticus** bacteriophages isolated from seawater. Appl. Environ. Microbiol. 44:466-470.

Remarks

History

History

Received from

Dr Tomio Kawata,
Department of Food Microbiology,
Tokushima University School of Medicine,
Tokushima,
770,
Japan

Date

02-21-1983

Isolated by

T. Koga and T. Kawata
Tokushima,
Japan

Date

06-1980

Source

Seawater

Updated at

2024-01-16