


<i>HER number</i> 169			
<b>Identification</b>			
<i>Name</i> 169	<i>Morphotype</i> B1 (Siphophage)	<i>Other designations</i>	
<b>Taxonomy</b>			
<i>Realm</i> <i>Duplodnaviria</i>	<i>Kingdom</i> <i>Heunggongvirae</i>	<i>Phylum</i> <u><i>Uroviricota</i></u>	<i>Class</i> <i>Caudoviricetes</i>
<i>Order</i>	<i>Family</i>	<i>Genus</i> <i>Enhodamvirus</i>	<i>Species</i> <i>Enhodamvirus VP5</i>
<b>Images</b>			
<i>Electron Micrograph</i> <i>Image</i> 	<i>Image description</i> Magnification: 297,000X Bar: 50 nm Staining: UA		
<i>H.-W.ACKERMAN III</i>			

<p><i>Characteristics</i></p> <p>Plaques of 0.5 mm, extremely opaque, more visible after 3 days. Elongated head.</p>	<p><i>Genomic sequence</i> Activated</p>
<h2>Propagation conditions</h2>	
<p><i>Bacterial hosts</i> 1169</p>	
<p><i>Reference</i></p> <p>Koga, T., S. Toyoshima, and T. Kawata. 1982. Morphological varieties and host ranges of <i>*Vibrio parahaemolyticus*</i> bacteriophages isolated from seawater. Appl. Environ. Microbiol. 44:466-470.</p>	
<p><i>Remarks</i></p>	
<h2>History</h2>	
<p><i>History</i></p> <p><b>Received from</b> Dr Tomio Kawata, Department of Food Microbiology, Tokushima University School of Medecine, Tokushima, 770, Japan</p>	<p><b>Date</b> 02-21-1983</p>
<p><b>Isolated by</b> T. Toga and T. Kawata Tokushima, Japan</p>	<p><b>Date</b> 07-1979</p>
<p><i>Source</i> Seawater</p>	
<p><i>Updated at</i> 2024-01-16</p>	