

<i>HER number</i> 191						
Identification						
<i>Name</i> 191	<i>Morphotype</i> B3 (Siphophage)	<i>Other designations</i> BLE				
Taxonomy						
<i>Realm</i> <i>Duplodnaviria</i>	<i>Kingdom</i> <i>Heunggongvira</i>	<i>Phylum</i> <i>Uroviricota</i>	<i>Class</i> <i>Caudoviricetes</i>			
<i>Order</i>	<i>Family</i>	<i>Genus</i>	<i>Species</i>			
Images						
<i>Electron Micrograph</i>	<i>Image description</i>					
<i>Characteristics</i> Plaques: 1.0 mm, clear.	<i>Genomic sequence</i> Deactivated					
Propagation conditions						
<i>Bacterial hosts</i> 1190						
<i>Reference</i> Doskocil, J., J. Forstova, and J. Stokrova. 1978. Temperate and virulent forms of phage θ attacking <i>*Bacillus licheniformis*</i> . Mol. Gen. Genet. 160:311-317.						
<i>Remarks</i> Phage BLE was isolated from the same batch independantly by Dr Ludvik: Ludvik, J., L. Erbenova, and H. Lipavska. 1977. Ultrastructure of <i>Bacillus licheniformis</i> bacteriophage BLE and its DNA. Virology 77:872-875.						

History

History

Received from

Dr. J. Doskocil

Institute of Molecular Genetics

Flemingovo nám. 2,

16637 Praha 6,

Czechoslovakia

Date

09-20-1983

Isolated by

J. Doskocil

Kourim,

Czechoslovakia

Date

1975

Source

A lysed industrial batch of **B. licheniformis** 10716.

Updated at

2024-01-16