

<p><i>HER number</i> 214</p>			
Identification			
<i>Name</i> 214	<i>Morphotype</i> A1 (Myophage)	<i>Other designations</i>	
Taxonomy			
<i>Realm</i>	<i>Kingdom</i>	<i>Phylum</i>	<i>Class</i>
<i>Order</i>	<i>Family</i> ?1	<i>Genus</i>	<i>Species</i>
Images			
<i>Electron Micrograph</i>		<i>Image description</i>	
<i>Characteristics</i> Plaques: 0.1mm, often smaller, opaque lysis only. Not chi-like. Contractile tail with ramified tail fibers.		<i>Genomic sequence</i> Deactivated	
Propagation conditions			
<i>Bacterial hosts</i> 1214			
<i>Reference</i> Rubires, X., S. Merino, A. Aguilar, M.M. Nogueras, and J.M. Tomas. 1998. Isolation of three different bacteriophages from mesophilic <i>*Aeromonas*</i> sp. that use different types of monopolar flagella as their primary receptor. FEMS Microbiol. Lett. 161:53-57.			

Remarks

History

History

Received from

Dr. Juan M. Thomas
Departament de Microbiologia,
Universitat de Barcelona,
Avgda. Diagonal 645,
08028 Barcelona,
Spain

Date

01-1998

Isolated by

Xavier Rubires
Departament de Microbiologia,
Universitat de Barcelona

Date

11-1994

Source

Lake water, Can Morgades lake, Barcelona, Spain.

Updated at

2024-01-17