

| | | | |
|--|--------------------------------------|--------------------------------------|----------------|
| <i>HER number</i> 349 | | | |
| Identification | | | |
| <i>Name</i> φ3T | <i>Morphotype</i> B1 (Siphophage) | <i>Other designations</i> phi3T | |
| Taxonomy | | | |
| <i>Realm</i> | <i>Kingdom</i> | <i>Phylum</i> | <i>Class</i> |
| <i>Order</i> | <i>Family</i> ?1 | <i>Genus</i> Spbetavirus | <i>Species</i> |
| Images | | | |
| <i>Electron Micrograph</i> | | <i>Image description</i> | |
| <i>Characteristics</i> Plaques: 0.5-1mm, clear. Typing phage of *B. subtilis* (No.6). | | <i>Genomic sequence</i> Activated | |
| Propagation conditions | | | |
| <i>Bacterial hosts</i> 1046 | | | |
| <i>Reference</i> Dean, D.H., C.L. Fort, J.A. Hoch. 1978. Characterization of temperate phages of *Bacillus subtilis*. Curr. Microbiol. 1: 213-217 | | | |

Remarks

History

History

Isolated by

R. Janvier

Félix d'Hérelle reference center for bacterial viruses
Québec, Canada

Date

02-02-1989

Received from

Dr. Donald H. Dean
Bacillus Genetic Stock Center
Dept. of Biochemistry
Ohio State University
484 W. 12th Ave, Columbus, Ohio 43210, USA

Date

15-09-1988

Source

Induction of *B. subtilis* 1L1 with mitomycin C

Updated at

2024-01-19