

Numéro HER
131

Identification

Nom Tb	Morphotype C1 (Podophage)	Autres désignations TB, Tbilisi	
-----------	------------------------------	------------------------------------	--


Taxonomie

Domaine <i>Duplodnaviria</i>	Règne <i>Heunggongvirae</i>	Embranchement Uroviricota	Classe <i>Caudoviricetes</i>
Ordre	Famille	Genre <i>Perisivirus</i>	Espèce <i>Perisivirus Tb</i>

Images

Photo en microscopie électronique

Image
FHRCEV



Description de l'image

Magnification: 297,000X

Bar: 50 nm

Staining: PTA

Caractéristique

Plaques: 0.5- 1mm, clear, sloping borders.
Host is class III pathogen.
Albimi broth, Serum dextrose agar.
Growth time = 36 h.

Séquence génomique

Activé

Conditions de propagation

Hôtes bactériens

1131

Référence

Brinley-Morgan, W. J. D. Kay, and D. E. Bradley. 1960. *Brucella* bacteriophage. Nature (London). 188:74-75.

Remarque

Historique

Historique

Received from

M. J. Corbel,
FAO/WHO Collaborating Centre for Reference and
Research on Brucellosis,
Ministry of Agriculture, Fisheries and Food,
Central Veterinary Laboratory,
New Haw, Weybridge, Surrey, England,
KT15 3NB

Date

06-22-1982

Isolated by

M. Z. Popkhadze, and T. G. Abashidze.
In Tiflis, Georgia,
URSS

Date

Source

Liquid manure of cowshed.

Dernière mise à jour

2024-01-16