Phage HER 154

Numéro HER
154

Identification


Nom	*Morphotype*	*Autres désignations*	
XP12	B2 (Siphophage)	Xp12, XP-12	

Taxonomie

Domaine	*Règne*	*Embranchement*	*Classe*
Duplodnaviria	*Heunggongvirae*	*Uroviricota*	*Caudoviricetes*
Ordre	*Famille*	*Genre*	*Espèce*

Images

Photo en microscopie électronique
Image



Description de l'image
Magnification: 297,000X
Bar: 50 nm
Staining: UAB

Caractéristique

Plaques: 2mm, clear.
Smaller plaques when cultivated without thymidine
Read after 24- 48 hrs.
Thymidine (5 mg/ml) needed in medium.
5-methylcytosine completely replaces cytosine.

Séquence génomique

Désactivé

Conditions de propagation

Hôtes bactériens

1154

Référence

Kuo, T. , T. Huang, and M. Teng. 1968. 5-methylcytosine replacing cytosine in the deoxyribonucleic acid of a bacteriophage for *Xanthomonas oryzae*. J. Mol. Biol. 34:373-375.

Remarque

Instructions for phage propagation and media composition are available upon request.
THIS PHAGE IS NOT AVAILABLE FOR THIS MOMENT

Historique

Historique

Received from
Dr Mélanie Ehrlich,
Department of Biochemistry,
Tulane University School of Medecine,
1430 Tulane Avenue,
New Orleans, LA, 70112,
USA.

Date
12-16-1982

Received from
Dr T. T. kuo,
Institute of Botany,
Academia Sinica,
Taiwan,
Republica of China.

Date

Isolated by
Kuo, Huang,
Wu and Cheng

Date
1968

Source
Water of a rice field, Taiwan

Dernière mise à jour
2024-01-16