

Numéro HER 191					
Identification					
<i>Nom</i> 191	<i>Morphotype</i> B3 (Siphophage)	<i>Autres désignations</i> BLE			
Taxonomie					
<i>Domaine</i>	<i>Règne</i>	<i>Embranchement</i>	<i>Classe</i>		
<i>Ordre</i>	<i>Famille</i> ?1	<i>Genre</i>	<i>Espèce</i>		
Images					
<i>Photo en microscopie électronique</i>		<i>Description de l'image</i>			
<i>Caractéristique</i> Plaques: 1.0 mm, clear.		<i>Séquence génomique</i> Désactivé			
Conditions de propagation					
<i>Hôtes bactériens</i> 1190					
<i>Référence</i>					
Doskocil, J., J. Forstova, and J. Stokrova. 1978. Temperate and virulent forms of phage θ attacking *Bacillus licheniformis*. Mol. Gen. Genet. 160:311-317.					
<i>Remarque</i>					
Phage BLE was isolated from the same batch independantly by Dr Ludvik: Ludvik, J., L. Erbenova, and H. Lipavska. 1977. Ultrastructure of *Bacillus licheniformis* bacteriophage BLE and its DNA. Virology 77:872-875.					
Historique					

Historique

Isolated by

J. Doskocil

Kourim,

Czechoslovakia

Date

1975

Received from

Dr. J. Doskocil

Institute of Molecular Genetics

Flemingovo nám. 2,

16637 Praha 6,

Czechoslovakia

Date

20-09-1983

Source

A lysed industrial batch of **B. licheniformis** 10716.

Dernière mise à jour

2024-01-16