Phage HER 191

Numéro HER
191

Identification

Nom	*Morphotype*	*Autres désignations*	
θc	B3 (Siphophage)	BLE	

Taxonomie

Domaine	*Règne*	*Embranchement*	*Classe*
Duplodnaviria	*Heunggongvirae*	*Uroviricota*	*Caudoviricetes*
Ordre	*Famille*	*Genre*	*Espèce*

Images

Photo en microscopie électronique	*Description de l'image*

Caractéristique	*Séquence génomique*
Plaques: 1.0 mm, clear.	Désactivé

Conditions de propagation

Hôtes bactériens
1190

Référence
Doskocil, J., J. Forstova, and J. Stokrova. 1978. Temperate and virulent forms of phage &theta ;attacking *Bacillus licheniformis*. Mol. Gen. Genet. 160:311-317.

Remarque

Phage BLE was isolated from the same batch independantly by Dr Ludvik:
Ludvik, J., L. Erbenova, and H. Lipavska. 1977. Ultrastructure of *Bacillus licheniformis* bacteriophage BLE and its DNA. Virology 77:872-875.

Historique

Historique

Received from
Dr. J. Doskocil
Institute of Molecular Genetics
Flemingovo nàm. 2,
16637 Praha 6,
Czechoslovakia

Date
09-20-1983

Isolated by
J. Doskocil
Kourim,
Czechoslovakia

Date
1975

Source
A lysed industrial batch of *B. licheniformis* 10716.

Dernière mise à jour
2024-01-16