

Numéro HER
317

Identification

<i>Nom</i> 317	<i>Morphotype</i> B1 (Siphophage)	<i>Autres désignations</i>	
-------------------	--------------------------------------	----------------------------	--

Taxonomie

<i>Domaine</i>	<i>Règne</i>	<i>Embranchement</i>	<i>Classe</i>
<i>Ordre</i>	<i>Famille</i> ?1	<i>Genre</i>	<i>Espèce</i>

Images

<i>Photo en microscopie électronique</i>	<i>Description de l'image</i>
<i>Caractéristique</i> Clear plaques of 1.0mm	<i>Séquence génomique</i> Désactivé

Conditions de propagation

<i>Hôtes bactériens</i> 1316

Référence

Jones, W.D. 1975. Differentiation of known strains of BCG from isolates of *Mycobacterium bovis* and *Mycobacterium tuberculosis* by using mycobacteriophage 33D. J. Clin. Microbiol. 1:391-392

Remarque

This phage lyses *M. tuberculosis* and wild strains of *M. bovis*; it does not lyse any of the attenuated strains of BCG.

Used in clinical settings for differentiation of BCG from *M. bovis* and *M. tuberculosis*. (1-RTD on ATCC 607)

Historique

Historique

Received from
Dr M. Buraczewska

Date

Received from
Dr Edith Mankiewicz
Lakeshore General Hospital
160 Stillview
Pointe-Claire, QC, Canada
H9R 2Y2

Date
14-04-1988

Isolated by
Dr M. Buraczewska
Watsaw, Poland

Date
1970

Source
Lysogenic bacilli

Dernière mise à jour
2024-01-19